Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

*Reports of States and yearly and monthly reports of cities of the United States.*

CALIFORNIA—*Los Angeles.*—Month of August, 1899. Estimated population, 103,000. Total number of deaths, 92, including enteric fever, 4; whooping cough, 1, and 4 from cerebro-spinal meningitis.

*Oakland.*—Month of August, 1899. Estimated population, 75,000. Total number of deaths, 80, including diphtheria, 1; enteric fever, 1, and 9 from phthisis pulmonalis.

*Sacramento.*—Month of August, 1899. Estimated population, 30,000. Total number of deaths, 23, including 4 from phthisis pulmonalis.

COLORADO—*Denver.*—Month of August, 1899. Estimated population, 170,000. Total number of deaths, 183, including diphtheria, 2; enteric fever, 10; whooping cough, 3; cerebro-spinal meningitis, 2, and 35 from phthisis pulmonalis.

CONNECTICUT.—Reports to the State board of health for the month of August, 1899, from 167 towns, having an aggregate estimated population of 911,524, show a total of 1,382 deaths, including diphtheria, 10; enteric fever, 21; measles, 1; scarlet fever, 4; whooping cough, 15; cerebro-spinal meningitis, 6, and 114 from phthisis pulmonalis.

*Hartford.*—Month of August, 1899. Census population, 53,230. Total number of deaths, 147, including diphtheria, 6; enteric fever, 2; scarlet fever, 1; whooping cough, 2, and 15 from phthisis pulmonalis.

IOWA—*Davenport.*—Month of August, 1899. Estimated population, 40,000. Total number of deaths, 40, including enteric fever, 1, and 6 from phthisis pulmonalis.

*Dubuque.*—Month of August, 1899. Estimated population, 45,000. Total number of deaths, 25, including diphtheria, 1; enteric fever, 2, and 3 from phthisis pulmonalis.

LOUISIANA—*New Orleans.*—Month of August, 1899. Estimated population, 300,000; white, 210,000; colored, 90,000. Total number of deaths, 532: white, 335; colored, 197, including enteric fever, 15; measles, 1; whooping cough, 2; cerebro-spinal meningitis, 3, and 59 from phthisis pulmonalis.

MASSACHUSETTS—*Brockton.*—Month of August, 1899. Estimated population, 37,278. Total number of deaths, 32, including spinal meningitis, 2, and 3 from phthisis pulmonalis.

*Fitchburg.*—Month of August, 1899. Estimated population, 29,000. Total number of deaths, 47, including diphtheria, 1; enteric fever, 1, and 4 from phthisis pulmonalis.

*Lawrence.*—Month of August, 1899. Estimated population, 58,500. Total number of deaths, 126, including diphtheria, 4; whooping 1, and 9 from phthisis pulmonalis.

MICHIGAN.—Reports to the State board of health, Lansing, for the week ended September 9, 1899, from 71 observers, indicate that intermittent fever decreased in area of prevalence. Phthisis pulmonalis was reported present at 199, enteric fever at 88, scarlet fever at 36,

diphtheria at 19, whooping cough at 15, measles at 13, cerebro-spinal meningitis at 2, and smallpox at 1 place—Maple Grove.

MINNESOTA—*Duluth.*—Month of May, 1899. Estimated population, 60,000. Total number of deaths, 45, including whooping cough, 1; cerebro-spinal meningitis, 1, and 4 from phthisis pulmonalis.

Month of June, 1899. Total number of deaths, 37, including enteric fever, 1, and 5 from phthisis pulmonalis.

Month of July, 1899. Total number af deaths, 29, including 1 from phthisis pulmonalis.

*St. Paul.*—Month of August, 1899. Estimated population, 200,000. Total number of deaths, 102, including diphtheria, 1; enteric fever, 3; scarlet fever, 1; whooping cough, 1, and 14 from phthisis pulmonalis.

NEW HAMPSHIRE—*Manchester.*—Month of August, 1899. Estimated population, 60,000. Total number of deaths, 97, including diphtheria, 1; whooping cough, 1, and 2 from phthisis pulmonalis.

NEW YORK—*Buffalo.*—Month of August, 1899. Estimated population, 370,000. Total number of deaths, 425, including diphtheria, 4; enteric fever, 14; measles, 1; whooping cough, 6; scarlet fever, 1; cerebro-spinal meningitis, 3, and 38 from phthisis pulmonalis.

OHIO—*Cleveland.*—Month of August, 1899. Estimated population, 395,000. Total number of deaths, 457, including diphtheria, 4; enteric fever, 10; scarlet fever, 1; whooping cough, 2; cerebro-spinal meningitis, 2, and 21 from phthisis pulmonalis.

*Toledo.*—Month of August, 1899. Estimated population, 150,000. Total number of deaths, 176, including diphtheria, 13; enteric fever, 5; scarlet fever, 3; whooping cough, 1, and 16 from phthisis pulmonalis.

TENNESSEE—*Chattanooga.*—Month of August, 1899. Estimated population, 40,000. Total number of deaths not reported. Five deaths from enteric fever.

TEXAS—*San Antonia.*—Month of August, 1899, Estimated population, 63,000. Total number of deaths, 125, including diphtheria, 1; enteric fever, 9; measles, 1; scarlet fever, 1; phthisis pulmonalis, 16, and 1 from smallpox.

VIRGINIA—*Petersburg.*—Month of July, 1899. Estimated population, 25,000. Total number of deaths, 57, including 4 from enteric fever.

Month of August. Estimated population, 25,000. Total number of deaths, 52, including enteric fever, 3, and 4 from phthisis pulmonalis.

WASHINGTON—*Seattle.*—Month of August, 1899. Estimated population, 75,000. Total number of deaths, 44, including diphtheria, 2; enteric fever, 3, and 2 from phthisis pulmonalis.

*Tacoma.*—Month of August, 1899. Estimated population, 50,000. Total number of deaths, 21, including diphtheria, 1, and 2 from phthisis pulmonalis.